



REFERENCES

- Andriani M, Drago G, Bernadi AM et al. 1995. Recombinant tissue plasminogen activator (rt-PA) as first-line therapy for declotting of haemodialysis access. *Nephrol Dial Transplant* 10: 1714-19.
- Antiplatelet Trialists Collaboration. 1994. Collaborative overview of randomised trials of antiplatelet therapy. II: Maintenance of vascular graft or arterial patency by antiplatelet therapy. *Br Med J* 308: 159-68.
- Ash SR, Mankus RA, Sutton JM et al. 1999. Concentrated sodium citrate as catheter lock solution. *J Am Soc Nephrol* 10: 272A.
- Ashleigh RJ, Al-Khaffaf H, Tronconi L et al. 1998. The use of metallic stents to treat central and peripheral venous stenosis related to haemodialysis access. *J Intervent Radiol* 13: 1-7.
- Atherikul K, Schwab SJ, Conlon PJ. 1998. Adequacy of haemodialysis with cuffed central venous catheters. *Nephrol Dial Transplant* 13: 745-49.
- Beathard GA. 1992. Percutaneous transvenous angioplasty in the treatment of vascular access stenosis. *Kidney Int* 42: 1390-97.
- Beathard GA. 1998. Physical examination of the dialysis vascular access. *Semin Dial* 11: 231-36.
- Beckingham IJ, O'Rourke JS, Bishop MC et al. 1993. Are back-up arteriovenous fistulae necessary for patients on continuous ambulatory peritoneal dialysis? *Lancet* 341: 1384-86.
- Bennion RS, Hiatt JR, Williams RA et al. 1985. A randomized, prospective study of perioperative antimicrobial prophylaxis for vascular access surgery. *J Cardiovasc Surg (Torino)* 26: 270-74.
- Besarab A, Samarapungam D. 1997. Measuring the adequacy of haemodialysis access. *Curr Opin Nephrol Hypertens* 5: 527-31.
- Bhatia DS, Money SR, Ochsner JL et al. 1996. Comparison of surgical bypass and percutaneous balloon dilatation with primary stent placement in the treatment of central venous obstruction in the dialysis patient: one-year follow-up. *Ann Vasc Surg* 10: 452-55.
- Boelaert JR, Van Landuyt HW, Godard CA et al. 1993. Nasal mupirocin ointment decreases the incidence of *Staphylococcus aureus* bacteraemias in haemodialysis patients. *Nephrol Dial Transplant* 8: 235-39.
- Buturovic J, Ponikvar R, Kandus A et al. 1998. Filling hemodialysis catheters in the interdialytic period: heparin versus citrate versus polygeline: a randomised study. *Artif Organs* 22: 945-47.
- Cayco AV, Abu-Alfa AK, Mahnensmith RL et al. 1998. Reduction in arteriovenous



- graft impairment. Results of a vascular access surveillance protocol. *Am J Kidney Dis* 32: 302-08.
- Cooper RI, Saad TF. 1999. Prevention of bacteremia in patients with tunneled cuffed permanent hemodialysis catheters using gentamicin catheter packing. *J Am Soc Nephrol* 10: (abstract 1032) 203A.
- Crain MR, Mewissen MW, Ostrowski GJ et al. 1996. Fibrin sleeve stripping for salvage of failing hemodialysis catheters: technique and initial results. *Radiology* 98: 41-44.
- Dapunt O, Feurstein M, Rendl KH et al. 1987. Transluminal angioplasty versus conventional operation in the treatment of haemodialysis fistula stenosis: results from a 5-year study. *Br J Surg* 74: 1004-05.
- Denys BG, Uretsky BF, Sudhakar R. 1993. Ultrasound-assisted cannulation of the internal jugular vein. A prospective comparison to the external landmark-guided technique. *Circulation* 87:1557-62.
- Farber A, Barbey MM, Grunert JH et al. 1999. Access-related venous stenoses and occlusions: treatment with percutaneous transluminal angioplasty and Dacron-covered stents. *Cardiovasc Intervent Radiol* 22: 214-18.
- Farrell J, Gellens M. 1997. Ultrasound guided cannulation versus the landmark-guided technique for acute haemodialysis access. *Nephrol Dial Transplant* 12: 1234-37.
- Farrell J, Walshe J, Gellens M et al. 1997. Complications associated with insertion of jugular venous catheters for haemodialysis: the value of postprocedural radiograph. *Am J Kidney Dis* 30: 690-92.
- Fong IW. 1993. Prevention of haemodialysis and peritoneal dialysis catheter related infection by topical povidone-iodine. *Postgrad Med J* 69 (Suppl 3): S15-17.
- Funaki B, Szymiski GX, Leef JA et al. 1997. Wallstent deployment to salvage dialysis graft thrombolysis complicated by venous rupture: early and intermediate results. *AJR Am J Roentgenol* 169: 1435-37.
- Glanz S, Gordon GH, Butt KM et al. 1987. The role of percutaneous angioplasty in the management of chronic hemodialysis fistulas. *Ann Surg* 206: 777-81.
- Gmelin E, Winterhoff R, Rinast E. 1989. Insufficient hemodialysis access fistulas: late results of treatment with percutaneous balloon angioplasty. *Radiology* 1989; 171: 657-60.
- Gualtieri E, Deppe SA, Sipperly MA et al. 1995. Subclavian venous catheterization. Greater success rate for less experienced operators using ultrasound guidance. *Crit Care Med* 23: 692-97.
- Hoehn B, Paul-Dauphin A, Hestin D et al. 1998. EPIBACDIAL: a multicenter prospective study of risk factors for bacteremia in chronic hemodialysis patients. *J Am Soc Nephrol* 9: 869-76.
- Kovalik EC, Newman GE, Suhocki P et al. 1994. Correction of central venous stenosis: use of angioplasty and vascular Wallstents. *Kidney Int* 45: 1177-81.
- Kovalik EC, Raymond JR, Albers FA. 1995. Catheter-related sepsis complicating long-term tunneled central venous dialysis catheters: management by guidewire
-



exchange. *Am J Kidney Dis* 25: 593-96.

Levin A, Mason AJ, Jindal KK et al. 1991. Prevention of hemodialysis subclavian vein catheter infections by topical povidone-iodine. *Kidney Int* 40: 934-38.

Lin BS, Kong CW, Tarrng DC et al. 1998. Anatomical variation of the internal jugular vein and its impact on temporary haemodialysis vascular access: an ultrasonographic survey in uraemic patients. *Nephrol Dial Transplant* 13: 134-38.

Lumsden AB, MacDonald MJ, Kikera D et al. 1998. Cost efficacy of duplex surveillance and prophylactic angioplasty of arteriovenous ePTFE. *Ann Vasc Surg* 12: 138-42. (No efficacy in reducing thrombosis with a 3-monthly prospective doppler screening protocol.)

Maki DG, Ringer M, Alvarado C. 1991. Prospective randomised trial of povidone-iodine, alcohol and chlorhexidine for the prevention of infection associated with central venous and arterial catheters. *Lancet* 338: 339-43.

Marr KA, Sexton DJ, Conlon PJ et al. 1997. Catheter-related bacteremia and outcome of attempted catheter salvage in patients undergoing hemodialysis. *Ann Intern Med* 127: 275-80.

Nadig C, Leidig M, Schmiedke T et al. 1998. The use of ultrasound for the placement of dialysis catheters. *Nephrol Dial Transplant* 13: 978-81.

Oakes DD, Sherck JP, Cobb LF. 1998. Surgical salvage of failed radiocephalic arteriovenous fistulae: techniques and results in 29 patients. *Kidney Int* 53: 480-87.

Patel N. 1998. Alternate approaches to central venous access. *Semin Intervent Radiol* 15: 325-33.

Patel RI, Peck SH, Cooper SG et al. 1998. Patency of Wallstents placed across the venous anastomosis of hemodialysis grafts after percutaneous recanalization. *Radiology* 209: 365-70.

Paulsen D, Reisotheer A, Aasen A et al. 1993. Use of tissue plasminogen activator for reopening of clotted dialysis catheters. *Nephron* 64: 468-70.

Paulson WD, Ram SJ, Birk CG et al. 1999. Does blood flow accurately predict thrombosis or failure of haemodialysis synthetic grafts? A meta-analysis. *Am J Kidney Dis* 34: 478-85.

Peacock SJ, Curtis N, Berendt AR et al 1999. Outcome following haemodialysis catheter-related *Staphylococcus aureus* bacteraemia. *J Hosp Infect* 41: 223-28.

Peska DN, DeLange B, Gratch JO et al. 1997. Short-term continuous infusion thrombolytic therapy for occluded central venous (sic) venous dialysis catheters. *Am J Manag Care* 3: 261-64.

Poulain F, Raynaud A, Bourquelot P et al. 1991. Local thrombolysis and thromboaspiration in the treatment of acutely thrombosed arteriovenous hemodialysis fistulas. *Cardiovasc Intervent Radiol* 14: 98-101.

Randolph AG, Cook DJ, Gonzales CA et al. 1996. Ultrasound guidance for placement of central venous catheters. A meta-analysis of the literature. *Crit Care Med* 24: 2053-58.

Roberts AB, Kahn MB, Bradford S et al. 1997. Graft surveillance and angioplasty



prolongs dialysis graft patency. *J Am Coll Surg* 183: 486-92.

Sands JJ, Patel S, Plaviak DJ et al. 1994. Pharmacomechanical thrombolysis with urokinase for treatment of thrombosed hemodialysis access grafts. *ASAIO J* 40: M886-88.

Schwab SJ, Raymond JR, Saeed M et al. 1989. Prevention of hemodialysis fistula thrombosis. Early detection of venous stenoses. *Kidney Int* 36: 707-11.

Sesso R, Barbosa D, Leme IL et al. 1998. Staphylococcus aureus prophylaxis in hemodialysis patients using central venous catheter: effect of mupirocin ointment. *J Am Soc Nephrol* 9: 1085-92.

Trerotola SO, Johnson MS, Schauwecker DS et al. 1996. Pulmonary emboli from pulse-spray and mechanical thrombolysis: evaluation with an animal dialysis-graft model. *Radiology* 200: 169-76.

Turmel-Rodrigues L, Pengloan J, Blanchier D et al. 1993. Insufficient dialysis shunts: improved long-term patency rates with close hemodynamic monitoring, repeated percutaneous balloon angioplasty, and stent placement. *Radiology* 187: 273-78.

Twardowski ZJ 1998. High dose intradialytic urokinase to restore the patency of permanent central venous hemodialysis catheters. *Am J Kidney Dis* 31: 841-47.

Wentworth DW, Kim FJ, Lentino JR et al. 1999. Citrate may be the anticoagulant of choice to prevent clotting of antibiotic blocks used in vascular access devices. *J Am Soc Nephrol* 10: (abstract 1127) 222A.

Winkler TA, Trerotola SO, Davidson DD et al. 1995. Study of thrombus from thrombosed hemodialysis access grafts. *Radiology* 197: 461-65.